

Recombinant Human NR1I3 293 Cell Lysate

Cat. No. NR1I3-3715HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for nuclear receptor subfamily 1, group I, member 3 (NR1I3), transcript variant 5 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

NR1I3 nuclear receptor subfamily 1, group I, member 3 [Homo sapiens]

Official Symbol

NR1I3

Synonyms

NR1I3; nuclear receptor subfamily 1, group I, member 3; nuclear receptor subfamily 1 group I member 3; CAR; CAR1; constitutive androstane receptor; MB67; constitutive active receptor; constitutive active response; orphan nuclear receptor MB67; orphan nuclear hormone receptor; constitutive activator of retinoid response; constitutive androstane nuclear receptor variant 2; constitutive androstane nuclear receptor variant 3; constitutive androstane nuclear receptor variant 4; constitutive androstane nuclear receptor variant 5; MGC97144; MGC97209; MGC150433;

Gene ID

9970

mRNA Refseq

NM_001077471

Protein Refseq

NP_001070939

MIM

603881

Chromosome Location

1q23.3

Pathway

Gene Expression, organism-specific biosystem; Generic Transcription Pathway, organism-specific biosystem; Glucocorticoid receptor regulatory network, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem; Nuclear Receptor transcription pathway, organism-specific biosystem; Nuclear Receptors, organism-specific biosystem; Nuclear receptors in lipid metabolism and toxicity, organism-specific biosystem;

Function

DNA binding; androgen receptor activity; ligand-activated sequence-specific DNA binding RNA polymerase II transcription factor activity; metal ion binding; receptor activity; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; steroid hormone receptor activity; thyroid hormone receptor activity; transcription coactivator activity; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA