

Recombinant Human NMT1 293 Cell Lysate

Cat. No. NMT1-3783HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for N-myristoyltransferase 1 (NMT1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[NMT1 N-myristoyltransferase 1 \[Homo sapiens \]](#)

Official Symbol

NMT1

Synonyms

NMT1; N-myristoyltransferase 1; glycylpeptide N-tetradecanoyltransferase 1; alternative; short form NMT S; long form; NMT L; myristoyl CoA:protein N myristoyltransferase; NMT; NMT 1; long form, NMT-L; alternative, short form NMT-S; type I N-myristoyltransferase; peptide N-myristoyltransferase 1; myristoyl-CoA:protein N-myristoyltransferase 1;

Gene ID

[4836](#)

mRNA Refseq

[NM_021079](#)

Protein Refseq

[NP_066565](#)

MIM

[160993](#)

UniProt ID

[P30419](#)

Chromosome Location

17q21

Pathway

Activation, myristoylation of BID and translocation to mitochondria, organism-specific biosystem; Apoptosis, organism-specific biosystem; Assembly of HIV virion,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem; Disease, organism-specific biosystem; HIV Infection, organism-specific biosystem; HIV Life Cycle, organism-specific biosystem; Intrinsic Pathway for Apoptosis, organism-specific biosystem;

Function

catalytic activity; glycylopeptide N-tetradecanoyltransferase activity; transferase activity, transferring acyl groups;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA