

Recombinant Human NFYC 293 Cell Lysate

Cat. No. NFYC-3838HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for nuclear transcription factor Y, gamma (NFYC), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

NFYC nuclear transcription factor Y, gamma [Homo sapiens]

Official Symbol

NFYC

Synonyms

NFYC; nuclear transcription factor Y, gamma; nuclear transcription factor Y subunit gamma; CBF C; NF YC; transactivator HSM-1; transactivator HSM-1/2; CCAAT binding factor subunit C; transcription factor NF-Y, C subunit; CAAT box DNA-binding protein subunit C; nuclear transcription factor Y subunit C; CCAAT transcription binding factor subunit gamma; histone H1 transcription factor large subunit 2A; HSM; CBFC; HAP5; CBF-C; NF-YC; H1TF2A; FLJ45775; DKFZp667G242;

Gene ID

4802

mRNA Refseq

NM_001142587

Protein Refseq

NP_001136059

MIM

605344

UniProt ID

Q13952

Chromosome Location

1p32

Pathway

Activation of Chaperone Genes by ATF6-alpha, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Activation of Chaperones by ATF6-alpha, organism-specific biosystem; Activation of Genes by ATF4, organism-specific biosystem; Antigen processing and presentation, organism-specific biosystem; Antigen processing and presentation, conserved biosystem; Diabetes pathways, organism-specific biosystem; Direct p53 effectors, organism-specific biosystem;

Function

DNA binding; protein binding; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; sequence-specific DNA binding transcription factor activity; transcription coactivator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA