

Recombinant Human NFE2 293 Cell Lysate

Cat. No. NFE2-3855HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for nuclear factor (erythroid-derived 2), 45kDa (NFE2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

NFE2 nuclear factor (erythroid-derived 2), 45kDa [Homo sapiens]

Official Symbol

NFE2

Synonyms

NFE2; nuclear factor (erythroid-derived 2), 45kDa; nuclear factor (erythroid derived 2), 45kD; transcription factor NF-E2 45 kDa subunit; NF E2; p45 NF-E2; leucine zipper protein NF-E2; nuclear factor, erythroid-derived 2 45 kDa subunit; p45; NF-E2;

Gene ID

4778

mRNA Refseq

NM_001136023

Protein Refseq

NP_001129495

MIM

601490

UniProt ID

Q16621

Chromosome Location

12q13

Pathway

Factors involved in megakaryocyte development and platelet production, organism-specific biosystem; Hemostasis, organism-specific biosystem;

Function

WW domain binding; WW domain binding; protein N-terminus binding; protein

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

dimerization activity; sequence-specific DNA binding; sequence-specific DNA binding
transcription factor activity; transcription coactivator activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA