

Recombinant Human NDUF7 293 Cell Lysate

Cat. No. NDUF7-3903HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 7, 18kDa (NDUF7), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

NDUFB7 NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 7, 18kDa [Homo sapiens]

Official Symbol

NDUFB7

Synonyms

NDUFB7; NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 7, 18kDa; NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 7 (18kD, B18); NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7; B18; CI B18; complex I B18 subunit; MGC2480; NADH ubiquinone oxidoreductase B18 subunit; complex I-B18; cell adhesion protein SQM1; NADH-ubiquinone oxidoreductase B18 subunit; CI-B18;

Gene ID

4713

mRNA Refseq

NM_004146

Protein Refseq

NP_004137

MIM

603842

UniProt ID

P17568

Chromosome Location

19p13.12

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Alzheimers disease, organism-specific biosystem; Alzheimers disease, conserved biosystem; Electron Transport Chain, organism-specific biosystem; Huntingtons disease, organism-specific biosystem; Huntingtons disease, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem;

Function

NADH dehydrogenase (ubiquinone) activity; NADH dehydrogenase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA