

Recombinant Human NDUFA8 293 Cell Lysate

Cat. No. NDUFA8-3915HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 8, 19kDa (NDUFA8), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

NDUFA8 NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 8, 19kDa [Homo sapiens]

Official Symbol

NDUFA8

Synonyms

NDUFA8; NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 8, 19kDa; NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 8 (19kD, PGIV); NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8; complex I PGIV subunit; MGC793; PGIV; complex I-19kD; complex I-PGIV; NADH:ubiquinone oxidoreductase PGIV subunit; NADH-ubiquinone oxidoreductase 19 kDa subunit; CI-19KD; CI-PGIV;

Gene ID

4702

mRNA Refseq

NM_014222

Protein Refseq

NP_055037

MIM

603359

UniProt ID

P51970

Chromosome Location

9q33.2

Pathway

Alzheimers disease, organism-specific biosystem; Alzheimers disease, conserved

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Electron Transport Chain, organism-specific biosystem; Huntingtons disease, organism-specific biosystem; Huntingtons disease, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem;

Function

NADH dehydrogenase (ubiquinone) activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA