

Recombinant Human NDUFA13 293 Cell Lysate

Cat. No. NDUFA13-3922HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 13 (NDUFA13), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

NDUFA13 NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 13 [Homo sapiens]

Official Symbol

NDUFA13

Synonyms

NDUFA13; NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 13; NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13; B16.6; CDA016; CGI 39; complex I B16.6 subunit; GRIM 19; GRIM19; CI-B16.6; complex I-B16.6; cell death-regulatory protein GRIM19; cell death regulatory protein GRIM-19; NADH-ubiquinone oxidoreductase B16.6 subunit; gene associated with retinoic and IFN-induced mortality 19 protein; gene associated with retinoic and interferon-induced mortality 19 protein; CGI-39; GRIM-19; FLJ58045; FLJ59191;

Gene ID

51079

mRNA Refseq

NM_015965

Protein Refseq

NP_057049

MIM

609435

UniProt ID

Q9P0J0

Chromosome Location

19p13.11

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Alzheimers disease, organism-specific biosystem; Alzheimers disease, conserved biosystem; EGFR1 Signaling Pathway, organism-specific biosystem; Huntingtons disease, organism-specific biosystem; Huntingtons disease, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem;

Function

ATP binding; NADH dehydrogenase (ubiquinone) activity; NADH dehydrogenase activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA