

Recombinant Human NCK1 293 Cell Lysate

Cat. No. NCK1-3947HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for NCK adaptor protein 1 (NCK1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[NCK1 NCK adaptor protein 1 \[Homo sapiens \]](#)

Official Symbol

NCK1

Synonyms

NCK1; NCK adaptor protein 1; NCK; cytoplasmic protein NCK1; NCKalpha; NCK tyrosine kinase; melanoma NCK protein; SH2/SH3 adaptor protein NCK-alpha; non-catalytic region of tyrosine kinase; nck-1; MGC12668;

Gene ID

[4690](#)

mRNA Refseq

[NM_001190796](#)

Protein Refseq

[NP_001177725](#)

MIM

[600508](#)

UniProt ID

[P16333](#)

Chromosome Location

3q21

Pathway

Activation of Rac, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Angiopoietin receptor Tie2-mediated signaling, organism-specific biosystem; Antigen Activates B Cell Receptor Leading to Generation of Second Messengers, organism-specific biosystem; Arf6 signaling events, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem;

Function

cytoskeletal adaptor activity; protein binding; protein domain specific binding; receptor binding; receptor signaling complex scaffold activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA