

Recombinant Human NAA50 293 Cell Lysate

Cat. No. NAA50-3991HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for N(alpha)-acetyltransferase 50, NatE catalytic subunit (NAA50) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

NAA50 N(alpha)-acetyltransferase 50, NatE catalytic subunit [Homo sapiens]

Official Symbol

NAA50

Synonyms

NAA50; N(alpha)-acetyltransferase 50, NatE catalytic subunit; MAK3, Mak3 homolog (S. cerevisiae) , N acetyltransferase 13 , N acetyltransferase 13 (GCN5 related) , NAT13; N-alpha-acetyltransferase 50; FLJ13194; NAT5; San; Mak3 homolog; separation anxiety; N-acetyltransferase 5; natE catalytic subunit; N-acetyltransferase san homolog; N-acetyltransferase 13 (GCN5-related); N-alpha-acetyltransferase 50, NatE catalytic subunit; SAN; MAK3; hSAN; NAT13; hNAT5;

Gene ID

80218

mRNA Refseq

NM_025146

Protein Refseq

NP_079422

MIM

610834

UniProt ID

Q9GZZ1

**Chromosome
Location**

3q13.31

Function

N-acetyltransferase activity; protein binding; transferase activity, transferring acyl

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



groups;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA