

Recombinant Human MUTYH 293 Cell Lysate

Cat. No. MUTYH-4052HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for mutY homolog (E. coli) (MUTYH), transcript variant beta3 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	MUTYH mutY homolog (E. coli) [Homo sapiens]
Official Symbol	MUTYH
Synonyms	MUTYH; mutY homolog (E. coli); mutY (E. coli) homolog; A/G-specific adenine DNA glycosylase; MYH; CYP2C; MGC4416;
Gene ID	4595
mRNA Refseq	NM_001048171
Protein Refseq	NP_001041636
MIM	604933
UniProt ID	Q9UIF7
Chromosome Location	1p34.1
Pathway	Base Excision Repair, organism-specific biosystem; Base excision repair, organism-specific biosystem; Base excision repair, conserved biosystem; Base-Excision Repair, AP Site Formation, organism-specific biosystem; Base-free sugar-phosphate removal via the single-nucleotide replacement pathway, organism-specific biosystem; Cleavage of the damaged purine, organism-specific biosystem; DNA Repair,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem;

Function

4 iron, 4 sulfur cluster binding; NOT MutLalpha complex binding; NOT MutLbeta complex binding; MutSalpha complex binding; NOT MutSbeta complex binding; catalytic activity; endonuclease activity; hydrolase activity, acting on glycosyl bonds; metal ion binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA