

Recombinant Human MPI 293 Cell Lysate

Cat. No. MPI-4235HCL Lot. No. (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for mannose phosphate isomerase (MPI) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[MPI mannose phosphate isomerase \[Homo sapiens \]](#)

Official Symbol

MPI

Synonyms

MPI; mannose phosphate isomerase; mannose-6-phosphate isomerase; mannose 6 phosphate isomerase; phosphohexomutase; phosphomannose isomerase 1; mannose-6- phosphate isomerase; PMI; PMI1; CDG1B; FLJ39201;

Gene ID

[4351](#)

mRNA Refseq

[NM_002435](#)

Protein Refseq

[NP_002426](#)

MIM

[154550](#)

UniProt ID

[P34949](#)

**Chromosome
Location**

15q22-qter

Pathway

Amino sugar and nucleotide sugar metabolism, organism-specific biosystem; Amino sugar and nucleotide sugar metabolism, conserved biosystem; Asparagine N-linked glycosylation, organism-specific biosystem; Biosynthesis of the N-glycan precursor (dolichol lipid-linked oligosaccharide, LLO) and transfer to a nascent protein,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem; D-mannose degradation, organism-specific biosystem;
Fructose and mannose metabolism, organism-specific biosystem; Fructose and
mannose metabolism, conserved biosystem;

Function

isomerase activity; mannose-6-phosphate isomerase activity; metal ion binding; zinc
ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA