

Recombinant Human MNAT1 293 Cell Lysate

Cat. No. MNAT1-4271HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for menage a trois homolog 1, cyclin H assembly factor (Xenopus laevis) (MNAT1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

MNAT1 menage a trois homolog 1, cyclin H assembly factor (*Xenopus laevis*) [*Homo sapiens*]

Official Symbol

MNAT1

Synonyms

MAT1; TFB3; CAP35; RNF66

Gene ID

4331

mRNA Refseq

NM_002431.3

Protein Refseq

NP_002422.1

MIM

602659

UniProt ID

P51948

Chromosome Location

14q23

Pathway

Basal transcription factors, organism-specific biosystem; Basal transcription factors, conserved biosystem; Cell Cycle, organism-specific biosystem; Cell Cycle, Mitotic, organism-specific biosystem; Cyclin A/B1 associated events during G2/M transition, organism-specific biosystem; Cyclin A:Cdk2-associated events at S phase entry, organism-specific biosystem; Cyclin D associated events in G1, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem;

Function

contributes_to DNA-dependent ATPase activity;contributes_to RNA polymerase II carboxy-terminal domain kinase activity;protein N-terminus binding;protein binding;contributes_to protein kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA