

Recombinant Mouse Arachidonate 12-lipoxygenase, His-tagged

Cat. No. Alox12-584M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Alox12 fused with N-terminal his-tagged protein, was expressed using the baculovirus system. MW = 76.4 kDa.
Species	Mouse
Source	Insect Cells
Description	12-Lipoxygenase(12-LO) catalyzes the formation of 12-HpETE from arachidonic acid. There are two main types of 12-LO, platelet-type and leukocyte-type, which share only 60% identity at the amino acid level and strongly differ in substrate specificity and enzyme kinetics. Leukocyte-type 12-LO has the ability to also produce 15-HpETE, however, platelet-type predominately forms 12-HpETE. Epidermal-type 12(S)- and 12(R)-LO enzymes have also been characterized, which catalyze the synthesis of their respective stereospecific 12-HETE products.
Purity	≥85%.
Formulation	50 mM sodium phosphate, pH 7.2, containing 100 mM sodium chloride and 20% glycerol.
Stability	≥6 months at -80°C; avoid freeze/thaw cycles by aliquoting the protein and storing at -80°C.

GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	Alox12 arachidonate 12-lipoxygenase [Mus musculus]
Synonyms	Alox12; arachidonate 12-lipoxygenase; P-12LO; Alox12p; 9930022G08Rik; arachidonate 12-lipoxygenase, 12S-type; 12S-LOX; 12S-lipoxygenase; OTTMUSP00000006289; platelet-type lipoxygenase 12; EC 1.13.11.31
Gene ID	11684
mRNA Refseq	NM_007440
Protein Refseq	NP_031466
UniProt ID	P39655
Chromosome Location	11 B3; 11 40.0 cM
Pathway	Arachidonic acid metabolism; Metabolic pathways
Function	arachidonate 12-lipoxygenase activity; iron ion binding; lipoxygenase activity; metal ion binding; oxidoreductase activity; oxidoreductase activity, acting on single donors with incorporation of molecular oxygen, incorporation of two atoms of oxygen