

Recombinant Human MED9 293 Cell Lysate

Cat. No. MED9-4377HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for mediator complex subunit 9 (MED9) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

MED9 mediator complex subunit 9 [Homo sapiens]

Official Symbol

MED9

Synonyms

MED9; mediator complex subunit 9; mediator of RNA polymerase II transcription, subunit 9 homolog (S. cerevisiae); mediator of RNA polymerase II transcription subunit 9; FLJ10193; MED25; mediator subunit 25; mediator of RNA polymerase II transcription, subunit 9 homolog; MGC138234;

Gene ID

55090

mRNA Refseq

NM_018019

Protein Refseq

NP_060489

MIM

609878

UniProt ID

Q9NWA0

Chromosome Location

17p11.2

Pathway

Developmental Biology, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

PPARA Activates Gene Expression, organism-specific biosystem; Regulation of Lipid Metabolism by Peroxisome proliferator-activated receptor alpha (PPARalpha), organism-specific biosystem; Transcriptional Regulation of White Adipocyte Differentiation, organism-specific biosystem;

Function

RNA polymerase II transcription cofactor activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA