

Recombinant Human MED4 293 Cell Lysate

Cat. No. MED4-4381HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for mediator complex subunit 4 (MED4) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

MED4 mediator complex subunit 4 [Homo sapiens]

Official Symbol

MED4

Synonyms

MED4; mediator complex subunit 4; mediator of RNA polymerase II transcription, subunit 4 homolog (S. cerevisiae) , VDRIP, vitamin D receptor interacting protein; mediator of RNA polymerase II transcription subunit 4; DRIP36; HSPC126; TRAP36; TRAP/SMCC/PC2 subunit p36 subunit; activator-recruited cofactor 36 kDa component; mediator of RNA polymerase II transcription, subunit 4 homolog; vitamin D3 receptor-interacting protein complex 36 kDa component; ARC36; VDRIP; RP11-90M2.2; FLJ10956;

Gene ID

29079

mRNA Refseq

NM_014166

Protein Refseq

NP_054885

MIM

605718

UniProt ID

Q9NPJ6

**Chromosome
Location**

13q14.12

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Developmental Biology, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Gene Expression, organism-specific biosystem; Generic Transcription Pathway, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins, organism-specific biosystem; PPARA Activates Gene Expression, organism-specific biosystem;

Function

RNA polymerase II transcription cofactor activity; RNA polymerase II transcription cofactor activity; ligand-dependent nuclear receptor transcription coactivator activity; protein binding; receptor activity; thyroid hormone receptor binding; transcription cofactor activity; vitamin D receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA