

Recombinant Human MED17 293 Cell Lysate

Cat. No. MED17-4391HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for mediator complex subunit 17 (MED17) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [MED17 mediator complex subunit 17 \[Homo sapiens \]](#)

Official Symbol MED17

Synonyms

MED17; mediator complex subunit 17; cofactor required for Sp1 transcriptional activation, subunit 6 (77kD) , cofactor required for Sp1 transcriptional activation, subunit 6, 77kDa , CRSP6; mediator of RNA polymerase II transcription subunit 17; CRSP77; DRIP80; TRAP80; ARC77; CRSP complex subunit 6; transcriptional coactivator CRSP77; activator-recruited cofactor 77 kDa component; vitamin D3 receptor-interacting protein complex 80 kDa component; thyroid hormone receptor-associated protein complex 80 kDa component; cofactor required for Sp1 transcriptional activation, subunit 6, 77kDa; CRSP6; FLJ10812;

Gene ID [9440](#)

mRNA Refseq [NM_004268](#)

Protein Refseq [NP_004259](#)

MIM [603810](#)

UniProt ID [Q9NVC6](#)

Chromosome Location 11q21

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Developmental Biology, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Gene Expression, organism-specific biosystem; Generic Transcription Pathway, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins, organism-specific biosystem; PPARA Activates Gene Expression, organism-specific biosystem;

Function

RNA polymerase II transcription cofactor activity; ligand-dependent nuclear receptor transcription coactivator activity; receptor activity; thyroid hormone receptor binding; transcription coactivator activity; transcription cofactor activity; vitamin D receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA