

Recombinant Human MED15 293 Cell Lysate

Cat. No. MED15-4392HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for mediator complex subunit 15 (MED15), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

MED15 mediator complex subunit 15 [Homo sapiens]

Official Symbol

MED15

Synonyms

MED15; mediator complex subunit 15; PC2 (positive cofactor 2, multiprotein complex) glutamine/Q rich associated protein , PCQAP, TNRC7, trinucleotide repeat containing 7; mediator of RNA polymerase II transcription subunit 15; Arc105; CAG7A; TIG 1; TPA inducible gene-1; CTG repeat protein 7a; TPA inducible protein; TPA-inducible gene 1 protein; trinucleotide repeat containing 7; activator-recruited cofactor, 105-kD; PC2-glutamine-rich-associated protein; PC2 glutamine/Q-rich-associated protein; activator-recruited cofactor 105 kDa component; trinucleotide repeat-containing gene 7 protein; positive cofactor 2 glutamine/Q-rich-associated protein; positive cofactor 2, glutamine/Q-rich-associated protein; PC2 (positive cofactor 2, multiprotein complex) glutamine/Q-rich-associated protein; TIG1; CTG7A; PCQAP; TIG-1; TNRC7; ARC105; FLJ42282; FLJ42935; DKFZp686A2214; DKFZp762B1216;

Gene ID

51586

mRNA Refseq

NM_001003891

Protein Refseq

NP_001003891

MIM

607372

UniProt ID

Q96RN5

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	22q11.2
Pathway	Developmental Biology, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Gene Expression, organism-specific biosystem; Generic Transcription Pathway, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins, organism-specific biosystem; PPARA Activates Gene Expression, organism-specific biosystem;
Function	RNA polymerase II transcription cofactor activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA