

Recombinant Human MECR 293 Cell Lysate

Cat. No. MECR-4394HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for mitochondrial trans-2-enoyl-CoA reductase (MECR), nuclear gene encoding mitochondrial protein, transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[MECR mitochondrial trans-2-enoyl-CoA reductase \[Homo sapiens \]](#)

Official Symbol

MECR

Synonyms

MECR; mitochondrial trans-2-enoyl-CoA reductase; trans-2-enoyl-CoA reductase, mitochondrial; CGI 63; FASN2B; mitochondrial 2 enoyl thioester reductase; NRBF1; nuclear receptor binding factor 1; NRBF-1; hsNrnf-1; nuclear receptor-binding factor 1; mitochondrial 2-enoyl thioester reductase; homolog of yeast 2-enoyl thioester reductase; CGI-63;

Gene ID

[51102](#)

mRNA Refseq

[NM_001024732](#)

Protein Refseq

[NP_001019903](#)

MIM

[608205](#)

UniProt ID

[Q9BV79](#)

Chromosome Location

1pter-p22.3

Pathway

Fatty Acid Biosynthesis, organism-specific biosystem; Fatty acid biosynthesis, elongation, mitochondria, organism-specific biosystem; Fatty acid biosynthesis,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

elongation, mitochondria, conserved biosystem; Fatty acid elongation, organism-specific biosystem; Fatty acid elongation, conserved biosystem; Metabolic pathways, organism-specific biosystem;

Function

nucleotide binding; oxidoreductase activity; receptor binding; trans-2-enoyl-CoA reductase (NADPH) activity; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA