

Recombinant Human MDM2 293 Cell Lysate

Cat. No. MDM2-4404HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for Mdm2 p53 binding protein homolog (mouse) (MDM2), transcript variant MDM2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[MDM2 Mdm2, p53 E3 ubiquitin protein ligase homolog \(mouse\) \[Homo sapiens \]](#)

Official Symbol

MDM2

Synonyms

MDM2; Mdm2, p53 E3 ubiquitin protein ligase homolog (mouse); Mdm2 p53 binding protein homolog (mouse) , Mdm2, transformed 3T3 cell double minute 2, p53 binding protein (mouse) , mouse double minute 2, human homolog of; p53 binding protein; E3 ubiquitin-protein ligase Mdm2; HDM2; HDMX; MGC5370; oncoprotein Mdm2; MDM2 variant FB28; MDM2 variant FB30; double minute 2, human homolog of; p53-binding protein; Mdm2, transformed 3T3 cell double minute 2, p53 binding protein; hdm2; ACTFS; MGC71221;

Gene ID

[4193](#)

mRNA Refseq

[NM_002392](#)

Protein Refseq

[NP_002383](#)

MIM

[164785](#)

UniProt ID

[Q00987](#)

**Chromosome
Location**

12q13-q14

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	AKT phosphorylates targets in the cytosol, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem; Apoptosis, organism-specific biosystem; Aurora A signaling, organism-specific biosystem; Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem;
Function	enzyme binding; ligase activity; metal ion binding; p53 binding; protein binding; ubiquitin-protein ligase activity; ubiquitin-protein ligase activity; zinc ion binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA