

Recombinant Human MDH2 293 Cell Lysate

Cat. No. MDH2-4407HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for malate dehydrogenase 2, NAD (mitochondrial) (MDH2), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name MDH2 malate dehydrogenase 2, NAD (mitochondrial) [Homo sapiens]

Official Symbol MDH2

Synonyms MDH2; malate dehydrogenase 2, NAD (mitochondrial); malate dehydrogenase, mitochondrial; MDH; MOR1; M-MDH; MGC:3559;

Gene ID 4191

mRNA Refseq NM_005918

Protein Refseq NP_005909

MIM 154100

UniProt ID P40926

Chromosome Location 7cen-q22

Pathway Citrate cycle (TCA cycle), organism-specific biosystem; Citrate cycle (TCA cycle), conserved biosystem; Citrate cycle, second carbon oxidation, 2-oxoglutarate =>oxaloacetate, organism-specific biosystem; Citrate cycle, second carbon oxidation, 2-oxoglutarate => oxaloacetate, conserved biosystem; Citric acid cycle (TCA cycle), organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

L-malate dehydrogenase activity; L-malate dehydrogenase activity; L-malate dehydrogenase activity; malate dehydrogenase (NADP+) activity; nucleotide binding; oxidoreductase activity; protein self-association;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA