

Recombinant Human MDH1 293 Cell Lysate

Cat. No. MDH1-4409HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for malate dehydrogenase 1, NAD (soluble) (MDH1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

MDH1 malate dehydrogenase 1, NAD (soluble) [Homo sapiens]

Official Symbol

MDH1

Synonyms

MDH1; malate dehydrogenase 1, NAD (soluble); malate dehydrogenase, cytoplasmic; soluble malate dehydrogenase; cytosolic malate dehydrogenase; MDHA; MOR2; MDH-s; MGC:1375;

Gene ID

4190

mRNA Refseq

NM_001199111

Protein Refseq

NP_001186040

MIM

154200

UniProt ID

P40925

**Chromosome
Location**

2p23

Pathway

Citrate cycle (TCA cycle), organism-specific biosystem; Citrate cycle (TCA cycle), conserved biosystem; Citrate cycle, second carbon oxidation, 2-oxoglutarate =>oxaloacetate, organism-specific biosystem; Citrate cycle, second carbon oxidation, 2-oxoglutarate => oxaloacetate, conserved biosystem; Gluconeogenesis, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem;

Function

L-malate dehydrogenase activity; malic enzyme activity; nucleotide binding;
oxidoreductase activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA