

Recombinant Human MCRS1 293 Cell Lysate

Cat. No. MCRS1-4411HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for microspherule protein 1 (MCRS1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	MCRS1 microspherule protein 1 [Homo sapiens]
Official Symbol	MCRS1
Synonyms	MCRS1; microspherule protein 1; ICP22BP; INO80 complex subunit Q; INO80Q; MCRS2; MSP58; P78; INO80 complex subunit J; 58 kDa microspherule protein; cell cycle-regulated factor p78; cell cycle-regulated factor (78 kDa);
Gene ID	10445
mRNA Refseq	NM_001012300
Protein Refseq	NP_001012300
MIM	609504
UniProt ID	Q96EZ8
Chromosome Location	12q13.12
Pathway	Herpes simplex infection, organism-specific biosystem; Herpes simplex infection, conserved biosystem;
Function	contributes_to histone acetyltransferase activity (H4-K16 specific); contributes_to

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

histone acetyltransferase activity (H4-K5 specific); contributes_to histone
acetyltransferase activity (H4-K8 specific); protein binding;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA