

Recombinant Human MAT2B 293 Cell Lysate

Cat. No. MAT2B-4453HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for methionine adenosyltransferase II, beta (MAT2B), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[MAT2B methionine adenosyltransferase II, beta \[Homo sapiens \]](#)

Official Symbol

MAT2B

Synonyms

MAT2B; methionine adenosyltransferase II, beta; methionine adenosyltransferase 2 subunit beta; MATIIBeta; SDR23E1; short chain dehydrogenase/reductase family 23E; member 1; MAT II beta; putative protein product of Nbla02999; dTDP-4-keto-6-deoxy-D-glucose 4-reductase; putative dTDP-4-keto-6-deoxy-D-glucose 4-reductase; short chain dehydrogenase/reductase family 23E, member 1; beta regulatory subunit of methionine adenosyltransferase; TGR; MAT-II; Nbla02999; MGC12237;

Gene ID

[27430](#)

mRNA Refseq

[NM_182796](#)

Protein Refseq

[NP_877725](#)

MIM

[605527](#)

UniProt ID

[Q9NZL9](#)

Chromosome Location

5q34-q35

Pathway

Biological oxidations, organism-specific biosystem; Cysteine and methionine

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

metabolism, organism-specific biosystem; Cysteine and methionine metabolism, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of amino acids and derivatives, organism-specific biosystem; Methionine degradation, organism-specific biosystem;

Function

dTDP-4-dehydrorhamnose reductase activity; methionine adenosyltransferase regulator activity; nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA