

Recombinant Human MALT1 293 Cell Lysate

Cat. No. MALT1-4527HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for mucosa associated lymphoid tissue lymphoma translocation gene 1 (MALT1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [MALT1 mucosa associated lymphoid tissue lymphoma translocation gene 1 \[Homo sapiens \]](#)

Official Symbol MALT1

Synonyms MALT1; mucosa associated lymphoid tissue lymphoma translocation gene 1; MLT; mucosa-associated lymphoid tissue lymphoma translocation protein 1; paracaspase; caspase-like protein; MALT associated translocation; MALT lymphoma-associated translocation; MALT-lymphoma associated translocation; MLT1; DKFZp434L132;

Gene ID [10892](#)

mRNA Refseq [NM_006785](#)

Protein Refseq [NP_006776](#)

MIM [604860](#)

UniProt ID [Q9UDY8](#)

Chromosome Location 18q21

Pathway Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; B cell receptor signaling pathway, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem; B cell receptor signaling pathway, conserved biosystem; BCR signaling pathway, organism-specific biosystem; Canonical NF-kappaB pathway, organism-specific biosystem; Downstream Signaling Events Of B Cell Receptor (BCR), organism-specific biosystem;

Function

cysteine-type endopeptidase activity; contributes_to kinase activator activity; peptidase activity; protein binding; protein self-association; signal transducer activity; ubiquitin-protein ligase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA