

Recombinant Human LMBRD1 293 Cell Lysate

Cat. No. LMBRD1-4716HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for LMBR1 domain containing 1 (LMBRD1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[LMBRD1 LMBR1 domain containing 1 \[Homo sapiens \]](#)

Official Symbol

LMBRD1

Synonyms

NESI; LMBD1; MAHCF; C6orf209; probable lysosomal cobalamin transporter; HDAG-L-interacting protein NESI; liver regeneration p-53 related protein; nuclear export signal-interacting protein; hepatitis delta antigen-L interacting protein

Gene ID

[55788](#)

mRNA Refseq

[NM_018368](#)

Protein Refseq

[NP_060838](#)

MIM

[612625](#)

UniProt ID

[Q9NUN5](#)

**Chromosome
Location**

6q13

Pathway

Defective BTB causes biotinidase deficiency, organism-specific biosystem; Defective MMADHC causes methylmalonic aciduria and homocystinuria type cblD, organism-specific biosystem; Metabolism, organism-specific biosystem

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

cobalamin binding

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA