

Recombinant Human LITAF 293 Cell Lysate

Cat. No. LITAF-4722HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for lipopolysaccharide-induced TNF factor (LITAF), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

LITAF lipopolysaccharide-induced TNF factor [Homo sapiens]

Official Symbol

LITAF

Synonyms

LITAF; lipopolysaccharide-induced TNF factor; lipopolysaccharide-induced tumor necrosis factor-alpha factor; FLJ38636; PIG7; SIMPLE; TP53I7; p53-induced gene 7 protein; LPS-induced TNF-alpha factor; tumor protein p53 inducible protein 7; lipopolysaccharide-induced TNF-alpha factor; small integral membrane protein of lysosome/late endosome; MGC116698; MGC116700; MGC116701; MGC125274; MGC125275; MGC125276;

Gene ID

9516

mRNA Refseq

NM_001136472

Protein Refseq

NP_001129944

MIM

603795

UniProt ID

Q99732

Chromosome Location

16p13.3-p12

Function

WW domain binding; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA