

Recombinant Human LDHA 293 Cell Lysate

Cat. No. LDHA-4790HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for lactate dehydrogenase A (LDHA), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

LDHA lactate dehydrogenase A [Homo sapiens]

Official Symbol

LDHA

Synonyms

LDHA; lactate dehydrogenase A; L-lactate dehydrogenase A chain; LDH-A; LDH-M; LDH muscle subunit; lactate dehydrogenase M; proliferation-inducing gene 19; renal carcinoma antigen NY-REN-59; cell proliferation-inducing gene 19 protein; LDH1; LDHM; GSD11; PIG19;

Gene ID

3939

mRNA Refseq

NM_001135239

Protein Refseq

NP_001128711

MIM

150000

UniProt ID

P00338

**Chromosome
Location**

11p15.1

Pathway

Cysteine and methionine metabolism, organism-specific biosystem; Cysteine and methionine metabolism, conserved biosystem; Glycolysis / Gluconeogenesis, organism-specific biosystem; Glycolysis / Gluconeogenesis, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Glycolysis and Gluconeogenesis, organism-specific biosystem; HIF-1-alpha transcription factor network, organism-specific biosystem; Metabolic pathways, organism-specific biosystem;

Function

L-lactate dehydrogenase activity; nucleotide binding; oxidoreductase activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA