

Recombinant Human LAT2 293 Cell Lysate

Cat. No. LAT2-4815HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for linker for activation of T cells family, member 2 (LAT2), transcript variant 3 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

LAT2 linker for activation of T cells family, member 2 [Homo sapiens]

Official Symbol

LAT2

Synonyms

LAT2; linker for activation of T cells family, member 2; WBSCR5, WBSCR15, Williams Beuren syndrome chromosome region 5; linker for activation of T-cells family member 2; HSPC046; LAB; linker for activation of B cells; linker for activation of T cells; transmembrane adaptor 2; non T cell activation linker; NTAL; WSCR5; non-T-cell activation linker; linker for activation of B-cells; membrane-associated adapter molecule; Williams-Beuren syndrome chromosomal region 5 protein; Williams-Beuren syndrome chromosomal region 15 protein; linker for activation of T cells, transmembrane adaptor 2; WBSCR5; WBSCR15;

Gene ID

7462

mRNA Refseq

NM_014146

Protein Refseq

NP_054865

MIM

605719

UniProt ID

Q9GZY6

**Chromosome
Location**

7q11.23

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	B Cell Receptor Signaling Pathway, organism-specific biosystem; I signaling in mast cells, organism-specific biosystem;
Function	SH2 domain binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA