

Recombinant Human KCNMB2 293 Cell Lysate

Cat. No. KCNMB2-5026HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for potassium large conductance calcium-activated channel, subfamily M, beta member 2 (KCNMB2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name KCNMB2 potassium large conductance calcium-activated channel, subfamily M, beta member 2 [Homo sapiens]

Official Symbol KCNMB2

Synonyms KCNMB2; potassium large conductance calcium-activated channel, subfamily M, beta member 2; calcium-activated potassium channel subunit beta-2; hbeta2; hbeta3; BKbeta2; hCG1646471; slo-beta-2; k(VCA)beta-2; BK channel subunit beta-2; MaxiK channel beta 2 subunit; maxi K channel subunit beta-2; charybdotoxin receptor subunit beta-2; large-conductance Ca²⁺-activated K⁺ channel beta2 subunit; calcium-activated potassium channel, subfamily M subunit beta-2; large conductance calcium-activated potassium channel beta 2 subunit; MGC22431;

Gene ID 10242

mRNA Refseq NM_181361

Protein Refseq NP_852006

MIM 605214

UniProt ID Q9Y691

Chromosome Location 3q26.32

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Ca²⁺ activated K⁺ channels, organism-specific biosystem; Hemostasis, organism-specific biosystem; Neuronal System, organism-specific biosystem; Nitric oxide stimulates guanylate cyclase, organism-specific biosystem; Platelet homeostasis, organism-specific biosystem; Potassium Channels, organism-specific biosystem; Vascular smooth muscle contraction, organism-specific biosystem;

Function

calcium-activated potassium channel activity; ion channel inhibitor activity; potassium channel activity; potassium channel regulator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA