

Recombinant Human Ectodysplasin A2 Receptor, Fc-tagged

Cat. No. EDA2R-2098H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Ectodysplasin A2 Receptor along with Fc tag is expressed in <i>HEK 293 cells</i> . The predicted molecular weight of Recombinant Human XEDAR is Mr 41.8 kDa. However, the actual molecular weight as observed by migration on SDS Page is Mr 61 kDa.
Species	Human
Source	HEK293
Description	Recombinant Human XEDAR containing the signal peptide from human CD33 joined with the extracellular domain of human XEDAR was fused to the Fc region of human IgG1 via a linker peptide. XEDAR (X-linked Ectodysplasin A2 Receptor) is a recently isolated member of the tumor necrosis factor receptor family that is highly expressed during embryonic development and induces apoptosis. XEDAR is a receptor for Ectodysplasin-A2 (EDA-A2) which may be a promising agent for the gene therapy of osteosarcoma.
Purity	>95% by SDS Page and analyzed by silver stain.
Biological Activity	The biological activity of Human XEDAR was determined by its ability to bind rhEDA-A2/His. Immobilized rhXEDAR/Fc at 1 µg/mL (100 µL/well) can bind rhEDA-A2/His with a linear range of 0.045 - 5 ng/mL in a functional ELISA.
Endotoxin	<1.0 EU/g as determined by the LAL method.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

State Of Matter	Lyophilized.
Formulation	This recombinant protein was 0.2 µm filtered and lyophilized from modified Dulbecco's phosphate buffered saline (1X PBS) pH 7.2 – 7.3 with no calcium, magnesium, or preservatives.
Storage And Stability	This lyophilized protein is stable for six to twelve months when stored desiccated at -20°C to -70°C. After aseptic reconstitution, this protein may be stored at 2°C to 8°C for one month or at -20°C to -70°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.

GENE INFORMATION

Gene Name	EDA2R ectodysplasin A2 receptor [Homo sapiens]
Synonyms	ectodysplasin A2 receptor; XEDAR; EDAA2R; EDA-A2R; TNFRSF27; EDA2R; X-linked ectodysplasin receptor; EDA-A2 receptor; X-linked ectodysplasin-A2 receptor; tumor necrosis factor receptor superfamily member XEDAR; OTTHUMP00000023448; ectodysplasin A2 receptor
Gene ID	60401
mRNA Refseq	NM_021783
Protein Refseq	NP_068555
MIM	300276
UniProt ID	Q9HAV5
Chromosome	Xq12

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Cytokine-cytokine receptor interaction
Function	protein binding; receptor activity; tumor necrosis factor receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA