

Recombinant Human IDH1 293 Cell Lysate

Cat. No. IDH1-5307HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for isocitrate dehydrogenase 1 (NADP+), soluble (IDH1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

IDH1 isocitrate dehydrogenase 1 (NADP+), soluble [Homo sapiens]

Official Symbol

IDH1

Synonyms

IDH1; isocitrate dehydrogenase 1 (NADP+), soluble; isocitrate dehydrogenase [NADP] cytoplasmic; NADP(+)-specific ICDH; oxalosuccinate decarboxylase; NADP-dependent isocitrate dehydrogenase, cytosolic; NADP-dependent isocitrate dehydrogenase, peroxisomal; IDH; IDP; IDCD; IDPC; PICD;

Gene ID

3417

mRNA Refseq

NM_005896

Protein Refseq

NP_005887

MIM

147700

UniProt ID

O75874

Chromosome Location

2q32-qter

Pathway

Citrate cycle (TCA cycle), organism-specific biosystem; Citrate cycle (TCA cycle), conserved biosystem; Citrate cycle, first carbon oxidation, oxaloacetate =>2-oxoglutarate, organism-specific biosystem; Citrate cycle, first carbon oxidation,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

oxaloacetate => 2-oxoglutarate, conserved biosystem; Glutathione metabolism, organism-specific biosystem;

Function

NAD binding; isocitrate dehydrogenase (NADP+) activity; isocitrate dehydrogenase (NADP+) activity; magnesium ion binding; oxidoreductase activity; oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor; protein homodimerization activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA