

Recombinant Human IBTK 293 Cell Lysate

Cat. No. IBTK-5317HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for inhibitor of Bruton agammaglobulinemia tyrosine kinase (IBTK) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

IBTK inhibitor of Bruton agammaglobulinemia tyrosine kinase [Homo sapiens]

Official Symbol

IBTK

Synonyms

IBTK; inhibitor of Bruton agammaglobulinemia tyrosine kinase; Bruton agammaglobulinemia tyrosine kinase inhibitor , BTKI; inhibitor of Bruton tyrosine kinase; DKFZP564B116; BTK-binding protein; inhibitor of Brutons tyrosine kinase; inhibitor of Brutons tyrosine kinase-beta; inhibitor of Brutons tyrosine kinase-alpha; inhibitor of Brutons tyrosine kinase-gamma; Bruton agammaglobulinemia tyrosine kinase inhibitor; BTKI; RP1-93K22.1; KIAA1417; MGC142256; MGC142258;

Gene ID

25998

mRNA Refseq

NM_015525

Protein Refseq

NP_056340

MIM

606457

UniProt ID

Q9P2D0

Chromosome Location

6q14.3

Pathway

BCR signaling pathway, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

protein kinase binding; protein tyrosine kinase inhibitor activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA