# Recombinant Human HSCB 293 Cell Lysate

**Cat. No.** HSCB-5382HCL **Lot. No.** (See product label)

## SPECIFICATION

| | |
|---|---|
| **Product Overview** | |
| **Species** | Human |
| **Source** | HEK293 |
| **Description** | Antigen standard for HscB iron-sulfur cluster co-chaperone homolog (E. coli) (HSCB) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection. |
| **Components** | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| **Size** | 0.1 mg |
| **Storage Instruction** | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment. |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| **Applications** | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane. |

## GENE INFORMATION

| | |
|---|---|
| **Gene Name** | HSCB HscB iron-sulfur cluster co-chaperone homolog (E. coli) [ Homo sapiens ] |
| **Official Symbol** | HSCB |
| **Synonyms** | HSCB; HscB iron-sulfur cluster co-chaperone homolog (E. coli); JAC1; HSC20; DNAJC20; dJ366L4.2; iron-sulfur cluster co-chaperone protein HscB, mitochondrial; J-type co-chaperone HSC20; dnaJ homolog subfamily C member 20; DnaJ (Hsp40) homolog, subfamily C, member 20; Jac1 |
| **Gene ID** | 150274 |
| **mRNA Refseq** | NM_172002 |
| **Protein Refseq** | NP_741999 |
| **MIM** | 608142 |
| **UniProt ID** | Q8IWL3 |
| **Chromosome Location** | 22q12.1 |
| **Pathway** | Metabolism of proteins, organism-specific biosystem; Mitochondrial Protein Import, organism-specific biosystem |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| **Function** | chaperone binding; heat shock protein binding; metal ion binding; molecular_function; protein binding |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA