

Recombinant Human HMMR 293 Cell Lysate

Cat. No. HMMR-5469HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for hyaluronan-mediated motility receptor (RHAMM) (HMMR), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

HMMR hyaluronan-mediated motility receptor (RHAMM) [Homo sapiens]

Official Symbol

HMMR

Synonyms

HMMR; hyaluronan-mediated motility receptor (RHAMM); hyaluronan mediated motility receptor; CD168; RHAMM; receptor for hyaluronan-mediated motility; intracellular hyaluronic acid-binding protein; IHABP; MGC119494; MGC119495;

Gene ID

3161

mRNA Refseq

NM_012484

Protein Refseq

NP_036616

MIM

600936

UniProt ID

O75330

Chromosome Location

5q33.2-qter

Pathway

ECM-receptor interaction, organism-specific biosystem; ECM-receptor interaction, conserved biosystem;

Function

hyaluronic acid binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA