

Recombinant Human HIST1H1B 293 Cell Lysate

Cat. No. HIST1H1B-5554HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for histone cluster 1, H1b (HIST1H1B) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[HIST1H1B histone cluster 1, H1b \[Homo sapiens \]](#)

Official Symbol

HIST1H1B

Synonyms

HIST1H1B; histone cluster 1, H1b; H1 histone family, member 5 , H1F5, histone 1, H1b; histone H1.5; H1.5; H1b; H1s 3; histone H1a; histone 1, H1b; H1 histone family, member 5; H1; H1B; H1F5; MGC126630; MGC126632;

Gene ID

[3009](#)

mRNA Refseq

[NM_005322](#)

Protein Refseq

[NP_005313](#)

MIM

[142711](#)

UniProt ID

[P16401](#)

**Chromosome
Location**

6p22.1

Pathway

Activation of DNA fragmentation factor, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptosis induced DNA fragmentation, organism-specific biosystem; Apoptotic executionphase, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

DNA binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA