

Recombinant Human HEXDC 293 Cell Lysate

Cat. No. HEXDC-5578HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for hexosaminidase (glycosyl hydrolase family 20, catalytic domain) containing (HEXDC) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

HEXDC hexosaminidase (glycosyl hydrolase family 20, catalytic domain) containing [Homo sapiens]

Official Symbol

HEXDC

Synonyms

HEXDC; hexosaminidase (glycosyl hydrolase family 20, catalytic domain) containing; hexosaminidase D; FLJ23825; beta-hexosaminidase D; beta-N-acetylhexosaminidase; N-acetyl-beta-galactosaminidase; hexosaminidase domain-containing protein;

Gene ID

284004

mRNA Refseq

NM_173620

Protein Refseq

NP_775891

UniProt ID

Q8WVB3

Chromosome Location

17q25.3

Pathway

Other glycan degradation, organism-specific biosystem; Other glycan degradation, conserved biosystem;

Function

beta-N-acetylhexosaminidase activity; cation binding; hydrolase activity, hydrolyzing

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

O-glycosyl compounds;

☎ Tel: 1-631-559-9269 1-516-512-3133

☎ Email: info@creative-biomart.com ☎ Fax: 1-631-938-8127

☎ 45-1 Ramsey Road, Shirley, NY 11967, USA