

Recombinant Human HDAC2 293 Cell Lysate

Cat. No. HDAC2-5605HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for histone deacetylase 2 (HDAC2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

HDAC2 histone deacetylase 2 [Homo sapiens]

Official Symbol

HDAC2

Synonyms

HDAC2; histone deacetylase 2; RPD3; YAF1; YY1-associated factor 1; transcriptional regulator homolog RPD3; HD2;

Gene ID

3066

mRNA Refseq

NM_001527

Protein Refseq

NP_001518

MIM

605164

UniProt ID

Q92769

**Chromosome
Location**

6q21

Pathway

Cell cycle, organism-specific biosystem; Cell cycle, organism-specific biosystem; Cell cycle, conserved biosystem; Chronic myeloid leukemia, organism-specific biosystem; Chronic myeloid leukemia, conserved biosystem; Delta-Notch Signaling Pathway, organism-specific biosystem; Direct p53 effectors, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

NAD-dependent histone deacetylase activity (H3-K14 specific); NAD-dependent histone deacetylase activity (H3-K9 specific); NAD-dependent histone deacetylase activity (H4-K16 specific); chromatin binding; enzyme binding; histone deacetylase activity; histone deacetylase activity (H3-K16 specific); hydrolase activity; protein binding; protein deacetylase activity; sequence-specific DNA binding; transcription factor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA