

Recombinant Human HAO2 293 Cell Lysate

Cat. No. HAO2-5637HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for hydroxyacid oxidase 2 (long chain) (HAO2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

HAO2 hydroxyacid oxidase 2 (long chain) [Homo sapiens]

Official Symbol

HAO2

Synonyms

HAO2; hydroxyacid oxidase 2 (long chain); hydroxyacid oxidase 2; (S) 2 hydroxy acid oxidase; GIG16; glycolate oxidase; growth inhibiting protein 16; HAOX2; long chain L 2 hydroxy acid oxidase; growth-inhibiting protein 16; long-chain L-2-hydroxy acid oxidase; long chain alpha-hydroxy acid oxidase; cell growth-inhibiting gene 16 protein; (S)-2-hydroxy-acid oxidase, peroxisomal;

Gene ID

51179

mRNA Refseq

NM_001005783

Protein Refseq

NP_001005783

MIM

605176

UniProt ID

Q9NYQ3

Chromosome Location

1p13.3-p13.1

Pathway

Glyoxylate and dicarboxylate metabolism, organism-specific biosystem; Glyoxylate and dicarboxylate metabolism, conserved biosystem; Metabolic pathways, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem; Peroxisome, organism-specific biosystem; Peroxisome, conserved biosystem;

Function

(S)-2-hydroxy-acid oxidase activity; FMN binding; long-chain-(S)-2-hydroxy-long-chain-acid oxidase activity; medium-chain-(S)-2-hydroxy-acid oxidase activity; oxidoreductase activity; very-long-chain-(S)-2-hydroxy-acid oxidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA