

Recombinant Human HADHB 293 Cell Lysate

Cat. No. HADHB-5645HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl-Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), beta subunit (HADHB), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

HADHB hydroxyacyl-CoA dehydrogenase/3-ketoacyl-CoA thiolase/enoyl-CoA hydratase (trifunctional protein), beta subunit [Homo sapiens]

Official Symbol

HADHB

Synonyms

HADHB; hydroxyacyl-CoA dehydrogenase/3-ketoacyl-CoA thiolase/enoyl-CoA hydratase (trifunctional protein), beta subunit; hydroxyacyl Coenzyme A dehydrogenase/3 ketoacyl Coenzyme A thiolase/enoyl Coenzyme A hydratase (trifunctional protein), beta subunit; trifunctional enzyme subunit beta, mitochondrial; mitochondrial trifunctional protein; beta subunit; MTPB; beta-ketothiolase; acetyl-CoA acyltransferase; 2-enoyl-Coenzyme A (CoA) hydratase, beta subunit; mitochondrial trifunctional enzyme, beta subunit; mitochondrial trifunctional protein, beta subunit; hydroxyacyl-Coenzyme A (CoA) dehydrogenase, beta subunit; 3-ketoacyl-Coenzyme A (CoA) thiolase of mitochondrial trifunctional protein, beta subunit; hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl-Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), beta subunit; ECHB; MSTP029; TP-BETA; MGC87480;

Gene ID

3032

mRNA Refseq

NM_000183

Protein Refseq

NP_000174

MIM

143450

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P55084
Chromosome Location	2p23
Pathway	Beta oxidation of decanoyl-CoA to octanoyl-CoA-CoA, organism-specific biosystem; Beta oxidation of hexanoyl-CoA to butanoyl-CoA, organism-specific biosystem; Beta oxidation of lauroyl-CoA to decanoyl-CoA-CoA, organism-specific biosystem; Beta oxidation of myristoyl-CoA to lauroyl-CoA, organism-specific biosystem; Beta oxidation of octanoyl-CoA to hexanoyl-CoA, organism-specific biosystem; Beta oxidation of palmitoyl-CoA to myristoyl-CoA, organism-specific biosystem; Fatty Acid Beta Oxidation, organism-specific biosystem;
Function	3-hydroxyacyl-CoA dehydrogenase activity; NAD binding; acetyl-CoA C-acyltransferase activity; enoyl-CoA hydratase activity; fatty-acyl-CoA binding; long-chain-3-hydroxyacyl-CoA dehydrogenase activity; long-chain-enoyl-CoA hydratase activity; protein binding; transferase activity, transferring acyl groups other than amino-acyl groups;