

Recombinant Human HADH 293 Cell Lysate

Cat. No. HADH-5646HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for hydroxyacyl-Coenzyme A dehydrogenase (HADH), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

HADH hydroxyacyl-CoA dehydrogenase [Homo sapiens]

Official Symbol

HADH

Synonyms

HADH; hydroxyacyl-CoA dehydrogenase; HADHSC, hydroxyacyl Coenzyme A dehydrogenase , L 3 hydroxyacyl Coenzyme A dehydrogenase, short chain; hydroxyacyl-coenzyme A dehydrogenase, mitochondrial; HADH1; SCHAD; short-chain 3-hydroxyacyl-CoA dehydrogenase; L-3-hydroxyacyl-Coenzyme A dehydrogenase, short chain; medium and short-chain L-3-hydroxyacyl-coenzyme A dehydrogenase; HAD; HCDH; HHF4; HADHSC; MSCHAD; MGC8392;

Gene ID

3033

mRNA Refseq

NM_001184705

Protein Refseq

NP_001171634

MIM

601609

UniProt ID

Q16836

Chromosome Location

4q22-q26

Pathway

Beta oxidation of butanoyl-CoA to acetyl-CoA, organism-specific biosystem; Beta

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

oxidation of decanoyl-CoA to octanoyl-CoA-CoA, organism-specific biosystem; Beta oxidation of hexanoyl-CoA to butanoyl-CoA, organism-specific biosystem; Beta oxidation of lauroyl-CoA to decanoyl-CoA-CoA, organism-specific biosystem; Beta oxidation of octanoyl-CoA to hexanoyl-CoA, organism-specific biosystem; Butanoate metabolism, organism-specific biosystem; Butanoate metabolism, conserved biosystem;

Function

3-hydroxyacyl-CoA dehydrogenase activity; NAD⁺ binding; coenzyme binding; nucleotide binding; oxidoreductase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA