

Recombinant Human GTF2F1 293 Cell Lysate

Cat. No. GTF2F1-5700HCL Lot. No. (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for general transcription factor IIF, polypeptide 1, 74kDa (GTF2F1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

GTF2F1 general transcription factor IIF, polypeptide 1, 74kDa [Homo sapiens]

Official Symbol

GTF2F1

Synonyms

GTF2F1; general transcription factor IIF, polypeptide 1, 74kDa; general transcription factor IIF, polypeptide 1 (74kD subunit); general transcription factor IIF subunit 1; BTF4; RAP74; TF2F1; TFIIF; TFIIF-alpha; transcription initiation factor RAP74; general transcription factor IIF 74 kDa subunit; transcription initiation factor IIF subunit alpha;

Gene ID

2962

mRNA Refseq

NM_002096

Protein Refseq

NP_002087

MIM

189968

UniProt ID

P35269

**Chromosome
Location**

19p13.3

Pathway

Abortive elongation of HIV-1 transcript in the absence of Tat, organism-specific biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Basal transcription factors, organism-specific biosystem; Basal transcription factors, conserved biosystem; Disease, organism-specific biosystem; Formation of HIV-1 elongation complex containing HIV-1 Tat, organism-specific biosystem; Formation of HIV-1 elongation complex in the absence of HIV-1 Tat, organism-specific biosystem;

Function

DNA binding; catalytic activity; phosphatase activator activity; protein binding; transcription coactivator activity; transcription factor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA