

Recombinant Human GNG2 293 Cell Lysate

Cat. No. GNG2-5853HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for guanine nucleotide binding protein (G protein), gamma 2 (GNG2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name GNG2 guanine nucleotide binding protein (G protein), gamma 2 [Homo sapiens]

Official Symbol GNG2

Synonyms GNG2; guanine nucleotide binding protein (G protein), gamma 2; guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2; g gamma-I; guanine nucleotide binding protein gamma 2; guanine nucleotide-binding protein G(I)/G(O) gamma-2 subunit;

Gene ID 54331

mRNA Refseq NM_001243773

Protein Refseq NP_001230702

MIM 606981

UniProt ID P59768

Chromosome Location 14q21

Pathway ADP signalling through P2Y purinoceptor 1, organism-specific biosystem; ADP signalling through P2Y purinoceptor 12, organism-specific biosystem; Activation of G protein gated Potassium channels, organism-specific biosystem; Activation of GABAB

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

receptors, organism-specific biosystem; Activation of Kainate Receptors upon glutamate binding, organism-specific biosystem; Aquaporin-mediated transport, organism-specific biosystem; CXCR3-mediated signaling events, organism-specific biosystem;

Function

signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA