

Recombinant Human GNG12 293 Cell Lysate

Cat. No. GNG12-5855HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for guanine nucleotide binding protein (G protein), gamma 12 (GNG12) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [GNG12 guanine nucleotide binding protein \(G protein\), gamma 12 \[Homo sapiens \]](#)

Official Symbol GNG12

Synonyms GNG12; guanine nucleotide binding protein (G protein), gamma 12; guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-12; G-protein gamma-12 subunit; FLJ31352; FLJ34695;

Gene ID [55970](#)

mRNA Refseq [NM_018841](#)

Protein Refseq [NP_061329](#)

UniProt ID [Q9UBI6](#)

Chromosome Location 1p31.2

Pathway

ADP signalling through P2Y purinoceptor 1, organism-specific biosystem; ADP signalling through P2Y purinoceptor 12, organism-specific biosystem; Activation of G protein gated Potassium channels, organism-specific biosystem; Activation of GABAB receptors, organism-specific biosystem; Activation of Kainate Receptors upon glutamate binding, organism-specific biosystem; Aquaporin-mediated transport, organism-specific biosystem; Calcium Regulation in the Cardiac Cell, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



specific biosystem;

Function

phosphate ion binding; signal transducer activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA