

Recombinant Human GCNT2 293 Cell Lysate

Cat. No. GCNT2-5979HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for glucosaminyl (N-acetyl) transferase 2, I-branching enzyme (I blood group) (GCNT2), transcript variant 3 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name GCNT2 glucosaminyl (N-acetyl) transferase 2, I-branching enzyme (I blood group) [Homo sapiens]

Official Symbol GCNT2

Synonyms GCNT2; glucosaminyl (N-acetyl) transferase 2, I-branching enzyme (I blood group); cataract, congenital , CCAT, GCNT5, glucosaminyl (N acetyl) transferase 2, I branching enzyme , glucosaminyl (N acetyl) transferase 2, I branching enzyme (li blood group) , glucosaminyl (N acetyl) transferase 5 , II, NACGT1; N-acetylglucosaminide beta-1,6-N-acetylglucosaminyl-transferase; bA360O19.2; bA421M1.1; IGNT; li blood group; NAGCT1; ULG3; unassigned linkage group 3; I beta-1,6-N-acetylglucosaminyltransferase; beta-1,6-N-acetylglucosaminyltransferase 2; II; CCAT; GCNT5; GCNT2C; NACGT1; MGC163396;

Gene ID 2651

mRNA Refseq NM_001491

Protein Refseq NP_001482

MIM 600429

UniProt ID Q06430

Chromosome 6p24.2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Glycosphingolipid biosynthesis - lacto and neolacto series, organism-specific biosystem; Glycosphingolipid biosynthesis - lacto and neolacto series, conserved biosystem; Metabolic pathways, organism-specific biosystem;
Function	N-acetyllactosaminide beta-1,6-N-acetylglucosaminyltransferase activity; transferase activity, transferring glycosyl groups;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA