

Recombinant Human FOSL1 293 Cell Lysate

Cat. No. FOSL1-6166HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for FOS-like antigen 1 (FOSL1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name FOSL1 FOS-like antigen 1 [Homo sapiens]

Official Symbol FOSL1

Synonyms FOSL1; FOS-like antigen 1; fos-related antigen 1; fra 1; FOS-like antigen-1; FRA; FRA1; fra-1;

Gene ID 8061

mRNA Refseq NM_005438

Protein Refseq NP_005429

MIM 136515

UniProt ID P15407

Chromosome Location 11q13

Pathway Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem; Calcium signaling in the CD4+ TCR pathway, organism-specific biosystem; DNA damage response (only ATM dependent), organism-specific biosystem; Downstream signaling in naive CD8+ T cells, organism-specific biosystem; HTLV-I infection, organism-specific biosystem; HTLV-I infection, conserved biosystem; Osteoclast differentiation, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

protein binding; protein dimerization activity; sequence-specific DNA binding;
sequence-specific DNA binding transcription factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA