

Recombinant Human FBXO6 293 Cell Lysate

Cat. No. FBXO6-6289HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for F-box protein 6 (FBXO6) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name FBXO6 F-box protein 6 [Homo sapiens]

Official Symbol FBXO6

Synonyms FBXO6; F-box protein 6; F box only protein 6; F-box only protein 6; FBG2; FBS2; FBX6; Fbx6b; F-box protein FBG2; F-box protein Fbx6; F-box/G-domain protein 2; F-box protein that recognizes sugar chains 2;

Gene ID 26270

mRNA Refseq NM_018438

Protein Refseq NP_060908

MIM 605647

UniProt ID Q9NRD1

Chromosome Location 1p36.22

Pathway Adaptive Immune System, organism-specific biosystem; Antigen processing: Ubiquitination and Proteasome degradation, organism-specific biosystem; Association of TriC/CCT with target proteins during biosynthesis, organism-specific biosystem; Chaperonin-mediated protein folding, organism-specific biosystem; Class I MHC mediated antigen processing & presentation, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

carbohydrate binding; glycoprotein binding; protein binding; ubiquitin-protein ligase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA