

Recombinant Human FAAH2 293 Cell Lysate

Cat. No. FAAH2-6480HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for fatty acid amide hydrolase 2 (FAAH2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

FAAH2 fatty acid amide hydrolase 2 [Homo sapiens]

Official Symbol

FAAH2

Synonyms

FAAH2; fatty acid amide hydrolase 2; AMDD, amidase domain containing; fatty-acid amide hydrolase 2; FAAH 2; FLJ31204; RP11 479E16.1; oleamide hydrolase 2; amidase domain containing; anandamide amidohydrolase 2; amidase domain-containing protein; AMDD;

Gene ID

158584

mRNA Refseq

NM_174912

Protein Refseq

NP_777572

MIM

300654

UniProt ID

Q6GMR7

Chromosome Location

Xp11.1

Pathway

anandamide degradation, organism-specific biosystem;

Function

carbon-nitrogen ligase activity, with glutamine as amido-N-donor; hydrolase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA