

Recombinant Human EXOSC5 293 Cell Lysate

Cat. No. EXOSC5-6500HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for exosome component 5 (EXOSC5) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

EXOSC5 exosome component 5 [Homo sapiens]

Official Symbol

EXOSC5

Synonyms

EXOSC5; exosome component 5; exosome complex component RRP46; exosome component Rrp46; hRrp46p; MGC12901; p12B; RRP41B; RRP46; Rrp46p; exosome complex exonuclease RRP46; ribosomal RNA-processing protein 46; chronic myelogenous leukemia tumor antigen 28; MGC111224;

Gene ID

56915

mRNA Refseq

NM_020158

Protein Refseq

NP_064543

MIM

606492

UniProt ID

Q9NQT4

Chromosome Location

19q13.1

Pathway

Activation of Genes by ATF4, organism-specific biosystem; Deadenylation-dependent mRNA decay, organism-specific biosystem; Destabilization of mRNA by Butyrate Response Factor 1 (BRF1), organism-specific biosystem; Destabilization of mRNA by

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

KSRP, organism-specific biosystem; Destabilization of mRNA by Tristetraprolin (TTP), organism-specific biosystem; Diabetes pathways, organism-specific biosystem; Disease, organism-specific biosystem;

Function

3-5-exoribonuclease activity; RNA binding; NOT exoribonuclease activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA