

Recombinant Human EXOSC3 293 Cell Lysate

Cat. No. EXOSC3-6502HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for exosome component 3 (EXOSC3), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

EXOSC3 exosome component 3 [Homo sapiens]

Official Symbol

EXOSC3

Synonyms

EXOSC3; exosome component 3; exosome complex component RRP40; CGI 102; CGI 102 protein; exosome component Rrp40; hRrp 40; hRrp40p; p10; RRP40; Rrp40p; exosome complex exonuclease RRP40; ribosomal RNA-processing protein 40; CGI-102; hRrp-40; bA3J10.7; MGC723; RP11-3J10.8; MGC15120;

Gene ID

51010

mRNA Refseq

NM_001002269

Protein Refseq

NP_001002269

MIM

606489

UniProt ID

Q9NQT5

Chromosome Location

9p11

Pathway

Activation of Genes by ATF4, organism-specific biosystem; Deadenylation-dependent mRNA decay, organism-specific biosystem; Destabilization of mRNA by Butyrate Response Factor 1 (BRF1), organism-specific biosystem; Destabilization of mRNA by

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

KSRP, organism-specific biosystem; Destabilization of mRNA by Tristetraprolin (TTP), organism-specific biosystem; Diabetes pathways, organism-specific biosystem; Disease, organism-specific biosystem;

Function

3-5-exoribonuclease activity; RNA binding; NOT exoribonuclease activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA