

Recombinant Human EXOSC2 293 Cell Lysate

Cat. No. EXOSC2-6503HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for exosome component 2 (EXOSC2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[EXOSC2](#) [exosome component 2](#) [[Homo sapiens](#)]

Official Symbol

EXOSC2

Synonyms

EXOSC2; exosome component 2; exosome complex component RRP4; homolog of yeast RRP4 (ribosomal RNA processing 4); 3 5 exoribonuclease (RRP4); hRrp4p; p7; RRP4; Rrp4p; exosome complex exonuclease RRP4; ribosomal RNA-processing protein 4; homolog of yeast RRP4 (ribosomal RNA processing 4), 3-5-exoribonuclease; homolog of yeast RRP4 (ribosomal RNA processing 4), 3 5 exoribonuclease (RRP4);

Gene ID

[23404](#)

mRNA Refseq

[NM_014285](#)

Protein Refseq

[NP_055100](#)

MIM

[602238](#)

UniProt ID

[Q13868](#)

Chromosome Location

9q34

Pathway

Activation of Genes by ATF4, organism-specific biosystem; Deadenylation-dependent

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA decay, organism-specific biosystem; Destabilization of mRNA by Butyrate Response Factor 1 (BRF1), organism-specific biosystem; Destabilization of mRNA by KSRP, organism-specific biosystem; Destabilization of mRNA by Tristetraprolin (TTP), organism-specific biosystem; Diabetes pathways, organism-specific biosystem; Disease, organism-specific biosystem;

Function

3-5-exoribonuclease activity; 7S RNA binding; NOT exoribonuclease activity; protein binding;